


§6.1 & §6.2 Anti-derivatives: Graphically, Numerically and Analytically

1. The graph of f is given in Figure 6.1. Please note that $f = F'$.

- What are the critical points of $F(x)$?
- Identify local extrema.
- Discuss concavity and points of inflection.

- d. Sketch an accurate graph of F , the anti-derivative of f . You are given that $F(0) = 0$. Fill in the chart


using the Fundamental Theorem of Calculus: $\int_a^b f(x)dx = F(b) - F(a)$. Then sketch F .



x	$\int_a^b f(x)dx$	$F(x)$
0	$\int_0^0 f(x)dx$	$0 + 0 = 0$
1	$\int_0^1 f(x)dx$	$0 + \quad =$
2		
3		
4		
5		

- e. On the same axes, now sketch F but given that $F(0) = 1$.


x	$F(x)$
0	1
1	
2	
3	
4	
5	



- f. Repeat again but now $F(0) = -1$

x	$F(x)$
0	-1
1	
2	
3	
4	
5	

2. Use the figure below and the fact that $P = 2$ when $t = 0$ to find values of P when $t = 1, 2, 3, 4$ and 5 .




x	$P(x)$
0	2
1	
2	
3	
4	
5	

3. Estimate $f(x)$ for $x = 2, 4, 6$, using the given values of $f'(x)$ and the fact that $f(0) = 50$.

x	0	2	4	6
$f'(x)$	17	15	10	2

4. A particle moves back and forth along the x-axis. The figure below (left one) approximates the velocity of the particle as a function of time. Positive velocities represent movement to the right and negative velocities represent movement to the left. The particle starts at the point $(5, 0)$. Graph the distance of the particle from the origin, with distance measured in kilometers and time in hours (LABEL!!!).



What is a function that has a derivative of $3x^2$?

A function is an **antiderivative** of f on an open interval I if $F'(x) = f(x)$ for all x in I.

There is an entire family of antiderivatives for each derivative. The **general** antiderivative of $3x^2$ is $x^3 + c$. c is called the **constant of integration**.

We use the following symbols: $\int f(x)dx = F(x) + c$

Constant of Integration
Antiderivative

Basic Rules for constructing Antiderivatives Analytically:

1. Zero

$$\int 0dx = c$$

2. Constant

$$\int k \, dx = kx + C$$

3. Power Rule

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

4. Natural Log

$$\int \frac{1}{x} dx = \ln |x| + C$$

5. Exponential

$$\int e^x dx = e^x + C$$

6. General Rules

$$\int kf(x)dx = k \int f(x)dx$$

$$\int (f(x) \pm g(x))dx = \int f(x)dx \pm \int g(x)dx$$

Example: Find the following:

a) $\int x^5 dx$

b) $\int x^{37} dx$

c) $\int x^{-2} dx$

d) $\int x^{\frac{1}{2}} dx$

e) $\int \frac{1}{x^3} dx$

f) $\int \sqrt[3]{x} dx$

g) $\int (2x^2 + 1)dx$

h) $\int (1 - x^3)dx$

i) $\int \left(\sqrt{x} - \frac{1}{\sqrt{x}} \right) dx$

j) $\int \sqrt{x}(x^2 + 1)dx$

k) $\int (x^2 + 1)(x + 2)dx$

l) $\int \left(\sqrt{x} + \frac{1}{2\sqrt{x}} \right) dx$

m) $\int \frac{x^3 - x^2 + 1}{x} dx$

n) $\int \frac{4x^5 - 3x^2 - 5}{x^2} dx$

o) $\int \frac{2x - x^2 + 3}{\sqrt{x}} dx$

p) $\int 8dx$

q) $\int 8xdx$

r) $\int 8x^2 dx$

s) $\int 8x^3 dx$

t) $\int \pi x^3 dx$

u) $\int \frac{e^x}{5} dx$

v) $\int (3x + x^2)dx$

w) $\int \frac{1}{2x} dx$

x) $\int \left(\frac{1}{x} + \frac{1}{x^3} \right) dx$

§6.1 – §6.2 Antiderivatives—cont.**Trig Rules:**

$$\begin{array}{ll} 1. \int \cos x dx = \sin x + c & 2. \int \sin x dx = -\cos x + c \\ 3. \int \sec^2 x dx = \tan x + c & 4. \int \sec x \tan x dx = \sec x + c \\ 5. \int \csc^2 x dx = -\cot x + c & 6. \int \csc x \cot x dx = -\csc x + c \end{array}$$

1. Find the following:

a) $\int (2 \cos x - \sin x) dx$

b) $\int 2 \sec x \tan x dx$

c) $\int -3 \csc^2 x dx$

d) $\int \frac{1}{2} \csc x \cot x dx$

e) $\int \sec x (\sec x + \tan x) dx$

f) $\int \frac{1}{\csc x} dx$

g) $\int (3x^2 - 5x + \sin x) dx$

h) $\int (e^x - \sec^2 x - \frac{5}{x}) dx$

i) $\int (\sin^2 x + \cos^2 x) dx$

j) $\int (10x^4 - 2 \sec^2 x) dx$

k) $\int \frac{\cos \theta}{\sin^2 \theta} d\theta$

2. Initial Conditions: You can solve for a particular c if you are given certain preliminary information; find $\int (2x + 3) dx$ such that $F(1) = 4$.

3. Find the velocity function $v(t)$ and position function $s(t)$ corresponding to the acceleration function $a(t) = 4t + 4$ given $v(0) = 8$ and $s(0) = 12$.

Highlight these three Rules:

$$1. \int a^x dx = \frac{a^x}{\ln a} + c$$

$$2. \int \frac{1}{\sqrt{1-x^2}} dx = \arcsin x + c$$

$$3. \int \frac{1}{1+x^2} dx = \arctan x + c$$

2. Find the following

$$a) \int (x^2 - 3x + 2^x) dx$$

$$b) \int \left(\frac{1}{3^{-x}} - x^3 \right) dx$$

$$c) \int \frac{2}{\sqrt{1-x^2}} dx$$

$$d) \int \left(\frac{1}{2x^2} - \frac{3}{1+x^2} \right) dx$$

$$e) \int \left(x^{\frac{2}{3}} - 4x^{\frac{-1}{5}} + \frac{4}{x} \right) dx$$

$$f) \int \left(\frac{2}{x} + 3e^x \right) dx$$

$$g) \int \left(\frac{1}{2t} - \sqrt{2}e^t \right) dt$$

$$h) \int (4 \sin x + 3 \cos x) dx$$

$$i) \int \left(\frac{1}{\csc x} \right) dx$$

$$j) \int (4 \sec^2 x + \csc x \cot x) dx$$

$$k) \int \sec x (\sec x + \tan x) dx$$

$$l) \int \frac{1-x+3\sqrt{x}}{x^2} dx$$

$$m) \int \sec x (\sec x + \cos x) dx$$

$$n) \int \frac{\sin x}{\cos^2 x} dx$$

$$o) \int \frac{\sin 2x}{\cos x} dx$$

$$p) \int \left(2^x - x^2 + \frac{1}{4^x} \right) dx$$

$$q) \int \left(e^x - \frac{1}{1+x^2} \right) dx$$

$$r) \int \left(3^x + \frac{1}{\sqrt{1-x^2}} \right) dx$$

§6.2—cont. Evaluating the Definite Integral

By the FTC part 1, we know the following: $\int_a^b f(x)dx = F(b) - F(a)$ where $f = F'$

Example 1: Evaluate $\int_1^5 2x dx$ by hand.

By FTC, we know $\int_1^5 2x dx = F(5) - F(1)$. Since $F(x) = \int 2x dx = x^2 + C$ we can conclude the following:

$$F(x) = x^2 + C$$

$$F(5) = 5^2 + C$$

$$F(5) - F(1) = (5^2 + C) - (1^2 + C) = 24$$

$$F(1) = 1^2 + C$$

Therefore $\int_1^5 2x dx = F(5) - F(1) = 24$

Example 2: Evaluate $\int_0^{\frac{5\pi}{6}} \cos x dx$ by hand. (*Notice the simpler notation*)

$$\int_0^{\frac{5\pi}{6}} \cos x dx = \sin x \Big|_0^{\frac{5\pi}{6}} = \sin\left(\frac{5\pi}{6}\right) - \sin(0) = \frac{1}{2} - 0 = \frac{1}{2} \quad \text{Use your calculator to check this answer.}$$

Practice: Evaluate the following using proper notation shown in Example 2. Check your answers with your calculator.

$$1. \int_0^1 x^3 dx$$

$$2. \int_{-2}^1 5 dx$$

$$3. \int_{-2}^4 \left(\frac{x}{2} + 3\right) dx$$

$$4. \int_{0.5}^{1.5} (-2x + 4) dx$$

$$5. \int_0^1 x dx$$

$$6. \int_0^2 (1 - x^2) dx$$


$$7. \int_0^1 e^x dx$$

$$8. \int_{-2}^1 |x| dx$$


Think Geometrically!

For each of the following problems, write a definite integral to express the area under the curve. The equation for the pictured function is provided. Then use the Fundamental Theorem of Calculus to find the area.


6. $f(x) = 2x + 1$




7. $f(x) = x^2 - 4$



8. $f(x) = 3 - x^2$




9. $f(x) = e^x$




For each of the following, graph and find the area under the graph of $f(x)$ from a to b . Show the definite integral and your evaluation.

10. $f(x) = 4 - x, \ a = -1, \ b = 2$



11. $f(x) = 4x - x^2, \ a = 0, \ b = 4$



12. $f(x) = \cos x, \ a = -\frac{\pi}{2}, \ b = \frac{\pi}{2}$

